

- 5 Dye Mixtures Of Fibre-Reactive Azo Dyes And Use Thereof For Dyeing
Material Containing Hydroxy- And/Or Carboxamido Groups


The present invention relates to the field of fibre-reactive dyes.

- 10 Dyestuffs containing chromophores linked via a piperazine type linking unit
are known from literature and are described for example in EP-A-0126265,
EP-A-0693538, WO99/05224 and WO00/08104.

- The inventors of the present invention have surprisingly found that mixtures
15 of dyestuffs of the general formula (I) and dyestuffs of the general formula
(II) give excellent application properties on cellulose containing material,
especially high levels of solubility in water or salt solution, high fixation
degrees, ease of washing out the unfixed dyestuff, good fastness to light
and water as well as robustness to process variables.

20


The present invention claims mixtures of fibre reactive dyes comprising one
or more dyestuffs of the formula (I)



25

and one or more dyestuffs of the general formula (II)

BEST AVAILABLE COPY



where

X^1 , X^2 independently are a labile atom or group;

5 Ar^1 is an aromatic residue substituted by at least one $-SO_3M$ group,
 M is hydrogen or alkali metal, especially sodium,


Ar^2 is an aromatic radical substituted with at least one $-SO_3M$ group,

a is 1 or 2

wherein,

10 if a is 2

L is a divalent radical typically of the form (y)



where

15 R^1 and R^2 are independently hydrogen, C_1 - C_4 alkyl optionally substituted by $-OR$, $-SR$, $-SO_3M$, or $-X$,

or a phenyl group optionally substituted by a sulfonic acid group, $-OR$, $-C_1$ - C_4 alkyl, or $-NR'COR$

20 L^1 is alkylene or arylene optionally substituted by a sulfonic acid group, $-OR$, $-C_1$ - C_4 alkyl or $-NR'COR$, $-COOR$

wherein R and R' are independently hydrogen or C_1 - C_4 alkyl and X is halogen

or R^1 and R^2 are independently optionally substituted alkyl, or

25 L is aminoethylpiperazine, under the proviso that if L is aminopiperazine, Ar^1 and Ar^2 are different or


if a is 1

L is a monovalent radical $-NR^3R^4$, $-SR^3$ or $-OR^3$

where

R^3 and R^4 have one of the meanings of R^1 and R^2 or for -

NR^3R^4 , R^3 and R^4 can form a cyclic structure of the form (o)



where

U is a C_4 - C_6 alkyl residue optionally substituted by a substituent of formula Z and optionally interrupted by heteroatoms or heteroatom-containing groups such as - O-, - NR^1 ,

n is 1, 2 or 3 and

Z is hydrogen, optionally substituted C_1 - C_4 alkyl, $-OR^5$, $-CO_2R^5$, $-COR^5$ and

R^5 is hydrogen, optionally substituted C_1 - C_4 alkyl, optionally substituted vinyl, optionally substituted phenyl.

(C_1 - C_4)-alkyl groups may be straight-chain or branched and are preferably for example methyl, ethyl, n-propyl, i-propyl or n-butyl. Substituted alkyl groups are preferably substituted by hydroxyl, (C_1 - C_4)-alkoxy, halogen or carboxyl groups. Substituted vinyl groups are for example $-C(CH_3)=CH_2$ or $-CH=CHCOOH$, substituted phenyl groups are for example phenyl substituted by $-COOH$, or $-SO_3M$.


Preferred embodiments of the invention are mixtures of one or more dyestuffs of the general formula (I) and one or more dyestuffs of the general formula (II)

where


X^1 and X^2 are independently chlorine, fluorine or 3 or 4-carboxypyridinium
especially preferred is chlorine,

Ar^1 and Ar^2 are independently a naphthyl residue substituted by at least one
sulfo group (q-1)

5




where n is 1 to 3, especially preferred (q-11) or (q-12)



(q-11)


or



(q-12)

10

or are a phenyl residue substituted by at least one sulfo group (q-2)



wherein

15


m is 1 or 2

p is 1 or 2

Y is independently hydrogen, halogen, R^5 , OR^5 , SR^5 , $NHCO R^5$,

where R^5 is as defined above, especially preferred Y is methyl.

20 When a is 2 L^1 is preferred to be an optionally substituted phenylene or an
alkylene residue optionally substituted or optionally interrupted by
heteroatoms or heteroatom containing groups, wherein L^1-N-R^1 or L^1-N-R^2
may contain a cyclic structural feature such as



where n and R^1 are as defined above and x is 2 to 5 and Z' has one of the meanings of Z .


When a is 1 L is preferably morpholine or N-Methylsulfanilic acid.

5

The dyestuffs of the formula (I) are contained in the mixture in quantity of 1% by weight to 99% by weight preferably in a mixing ratio of 10% by weight to 90% by weight and the dyestuffs of the formula (II) are contained in the mixture in a mixing ratio of 99% by weight to 1% by weight, preferably in a mixing ratio of 90% by weight to 10% by weight.

10

Dyestuff mixtures according to the invention can be obtained upon reacting chromophores of formula (III)



(III)

15

with an appropriate mixture of 2-aminoethylpiperazine and a diamine H-L-H, or amine H-L, wherein L is as defined above, followed by precipitation using methylated spirits and conventional filtration.

20

The dyestuffs of the present invention can be present as a preparation in solid or liquid (dissolved) form. In solid form they generally contain the electrolyte salts customary in the case of water-soluble and in particular fibre-reactive dyes, such as sodium chloride, potassium chloride and sodium

- sulfate, and also the auxiliaries customary in commercial dyes, such as buffer substances capable of establishing a pH in aqueous solution between 3 and 8, such as sodium acetate, sodium borate, sodium bicarbonate, sodium citrate, sodium dihydrogenphosphate and disodium
- 5 hydrogenphosphate, small amounts of siccatives or, if they are present in liquid, aqueous solution (including the presence of thickeners of the type customary in print pastes), substances which ensure the permanence of these preparations, for example mold preventatives.
- 10 In general, the dyestuff mixtures of the present invention are present as dye powders containing 10 to 80% by weight, based on the dye powder or preparation, of a strength-standardizing colorless diluent electrolyte salt, such as those mentioned above. These dye powders may additionally include the aforementioned buffer substances in a total amount of up to 10%, based
- 15 on the dye powder. If the dye mixtures of the present invention are present in aqueous solution, the total dye content of these aqueous solutions is up to about 50 % by weight, for example between 5 and 50% by weight, and the electrolyte salt content of these aqueous solutions will preferably be below 10% by weight, based on the aqueous solutions. The aqueous solutions
- 20 (liquid preparations) may include the aforementioned buffer substances in an amount which is generally up to 10% by weight, for example 0.1 to 10% by weight, preference being given to up to 4% by weight, especially 2 to 4% by weight.
- 25 The dyestuff mixtures of the instant invention are reactive dyestuffs suitable for dyeing and printing hydroxy- and/or carboxamido-containing fibre materials by the application and fixing methods numerously described in the art for fibre-reactive dyes. They provide exceptionally strong and economic shades. Such dyes especially when used for exhaust dyeing of cellulosic
- 30 materials can exhibit excellent properties including build-up, light-fastness, high levels of solubility in water or salt solution, high fixation degrees, ease of washing out the unfixed dyestuff, as well as robustness to process

variables.

They are also wholly compatible with similar dyes designed for high temperature (80-100°C) application to cellulosic textiles, and thus lead to highly reproducible application processes, with short application times. The present invention therefore also provides for use of the inventive dyestuffs for dyeing and printing hydroxy- and/or carboxamido-containing fibre materials and processes for dyeing and printing such materials using a dyestuff according to the invention. Usually the dyestuff is applied to the substrate in dissolved form and fixed on the fibre by the action of an alkali or by heating or both.

Hydroxy-containing materials are natural or synthetic hydroxy-containing materials, for example cellulose fiber materials, including in the form of paper, or their regenerated products and polyvinyl alcohols. Cellulose fiber materials are preferably cotton but also other natural vegetable fibers, such as linen, hemp, jute and ramie fibres. Regenerated cellulose fibers are for example staple viscose and filament viscose.

Carboxamido-containing materials are for example synthetic and natural polyamides and polyurethanes, in particular in the form of fibers, for example wool and other animal hairs, silk, leather, nylon-6,6, nylon-6, nylon-11, and nylon-4.

Application of the inventive dyestuffs is by generally known processes for dyeing and printing fiber materials by the known application techniques for fibre-reactive dyes. The dyestuffs according to the invention are highly compatible with similar dyes designed for high temperature (80-100°C) applications and are advantageously useful in exhaust dyeing processes.

30

Similarly, the conventional printing processes for cellulose fibers, which can either be carried out in single-phase, for example by printing with a print

paste containing sodium bicarbonate or some other acid-binding agent and the colorant, and subsequent steaming at appropriate temperatures, or in two phases, for example by printing with a neutral or weakly acid print paste containing the colorant and subsequent fixation either by passing the printed material through a hot electrolyte-containing alkaline bath or by overpadding with an alkaline electrolyte-containing padding liquor and subsequent batching of this treated material or subsequent steaming or subsequent treatment with dry heat, produce strong prints with well defined contours and a clear white ground. Changing fixing conditions has only little effect on the outcome of the prints. Not only in dyeing but also in printing the degrees of fixation obtained with dye mixtures of the invention are very high. The hot air used in dry heat fixing by the customary thermofix processes has a temperature of from 120 to 200°C. In addition to the customary steam at from 101 to 103°C, it is also possible to use superheated steam and high pressure steam at up to 160°C.

The inventive dyestuffs can in addition be used to produce inks useful for printing the substrates described above, for example textiles, especially cellulosic textiles, and paper. Such inks can be used in all technologies, for example conventional printing, ink-jet printing or bubble-jet printing (for information on such printing technologies see for example Text. Chem. Color, Volume 19(8), pages 23 ff and Volume 21, pages 27 ff).

Acid-binding agents responsible for fixing the dyes to cellulose fibers are for example water-soluble basic salts of alkali metals and of alkaline earth metals of inorganic or organic acids, and compounds, which release alkali when hot. Of particular suitability are the alkali metal hydroxides and alkali metal salts of weak to medium inorganic or organic acids; the preferred alkali metal compounds being the sodium and potassium compounds. These acid-binding agents are for example sodium hydroxide, potassium hydroxide, sodium carbonate, sodium bicarbonate, potassium carbonate, sodium formate, sodium dihydrogenphosphate and disodium hydrogenphosphate.

Treating the dyestuffs according to the invention with the acid-binding agents, with or without heating, bonds the dyes chemically to the cellulose fibers. Especially the dyeings on cellulose, after they have been given the usual aftertreatment of rinsing to remove unfixed dye portions, show excellent properties.

The dyeings of polyurethane and polyamide fibers are customarily carried out from an acid medium. The dyebath may contain for example acetic acid and/or ammonium sulfate and/or acetic acid and ammonium acetate or sodium acetate to bring it to the desired pH. To obtain a dyeing of acceptable levelness it is advisable to add customary leveling auxiliaries, for example based on a reaction product of cyanuric chloride with three times the molar amount of an aminobenzenesulfonic acid or aminonaphthalenesulfonic acid or based on a reaction product of for example stearylamine with ethylene oxide. In general the material to be dyed is introduced into the bath at a temperature of about 40°C and agitated therein for some time, the dyebath is then adjusted to the desired weakly acid, preferably weakly acetic acid, pH, and the actual dyeing is carried out at temperature between 60 and 98°C. However, the dyeings can also be carried out at the boil or at temperatures up to 120°C (under superatmospheric pressure).


The examples hereinbelow serve to illustrate the invention. Parts and percentages are by weight, unless otherwise stated. Parts by weight relate to parts by volume as the kilogram relates to the liter. The compounds described in the examples in terms of a formula are indicated in the form of the free sulphonic acids, but as in general they are prepared and isolated in the form of their alkali metal salts, such as lithium, sodium or potassium salts, and used for dyeing in the form of these salts. The starting compounds and components mentioned in the form of the free acid in the examples hereinbelow may be used in the synthesis as such or similarly in the form of

their salts, preferably alkali metal salts.


5 Example 1

A 1:1 molar mixture of 2-aminoethylpiperazine (4) and ethylenediamine (5) (0.006 mol) was added to a stirred suspension of the dichlorotriazinyl red dye(3) (0.012 mol) in water (350 mols) at pH 6. The pH was adjusted and maintained at pH 10 by the addition of 2N NaOH solution. After three
 10 hours, chromatography indicated that the reaction was complete and the pH was adjusted to 6 with 2N HCl before precipitating the product by addition of methylated spirits. The resulting solid was filtered off and dried to give a dark red powder (11.3g). Analytical data were consistent with a 1:1 mixture of the two dyes (6) and (7).

15




The examples herein below describe further dye mixtures according to the invention. They may be prepared according to the invention either by mechanically mixing the individual dyes or else chemically, for example similarly to the above illustrative embodiment, using as starting compounds of general formula (III), in which X is for example chlorine and Ar¹ and Ar² are for example:





10

2-aminopiperazine and a diamine H-L-H or amine H-L, wherein L is for example a residue of formula a to af



Following exactly analogous procedures the following dyes (examples 2 – 56) were synthesized giving 1:1 mixtures of the respective dyestuff (8) and

5 (9)




Example No.	Ar ¹	Ar ²	L
2	A	A	a
3	A	A	b
4	A	A	c
5	A	A	d
6	A	A	e
7	A	A	f
8	A	A	g
9	A	A	h
10	A	A	k
11	A	A	l

Example No.	Ar ¹	Ar ²	L
12	A	A	p
13	B	B	a
14	B	B	d
15	B	B	g
16	B	B	k
17	B	B	l
18	B	B	m
19	B	B	o
20	B	B	p
21	A	G	j
22	A	G	k
23	A	G	l
24	B	F	k
25	B	F	l
26	B	F	o
27	B	F	p
28	C	C	a
29	C	C	k
30	C	C	l
31	F	F	a
32	F	F	k
33	F	F	l
34	D	D	k
35	D	D	o
36	E	E	l
37	H	H	m
38	H	H	n
39	G	G	l

Example No.	Ar ¹	Ar ²	L
40	G	G	n
41	B	B	v
42	F	F	v
43	B	B	w
44	F	F	w
45	B	B	x
46	F	F	x
47	B	B	f
48	F	F	y
49	F	F	z
50	B	B	aa
51	B	B	j
52	F	F	j
53	F	F	o
54	A	A	ab
55	B	B	ae
56	B	B	af

Examples 57-71 consist of mixtures of dyes of the form (8) and (10).




Example No.	Ar ¹	Ar ²	L
57	A	A	q
58	A	A	r
59	A	A	u
60	A	A	t
61	B	B	q
62	B	B	s
63	B	B	t
64	B	B	u
65	F	F	q
66	F	F	s
67	F	F	t
68	C	C	s
69	C	C	t
70	G	G	u
71	B	B	ad

All these dyestuff mixtures give excellent application properties on cellulose containing material, especially high levels of solubility in water or salt solution, high fixation degrees, ease of washing out the unfixed dyestuff, good fastness to light and water as well as robustness to process variables.


What is claimed is:

1. Mixture of fibre reactive dyes comprising one or more dyestuffs of the formula (I)

5



and one or more dyestuffs of the general formula (II)



10

where

X^1 , X^2 are independently a labile atom or group;

Ar^1 is an aromatic residue substituted by at least one $-SO_3M$ group;

M is hydrogen or alkali metal, especially sodium;

15 Ar^2 is an aromatic radical substituted with at least one $-SO_3M$ group;


a is 1 or 2

wherein,

if a is 2

L is a divalent radical typically of the form (y)

20



where

R^1 and R^2 are independently hydrogen, C_1 - C_4 alkyl optionally substituted by -OR, -SR, - SO_3M or X, or a phenyl group optionally substituted by a sulfonic acid group, -OR, - C_1 - C_4 -alkyl, or $NR'COR$ and

L^1 is arylene or alkylene optionally substituted by a sulfonic acid group, -OR, - C_1 - C_4 -alkyl - COOR, or $NR'COR$

or


L is aminoethylpiperazine, under the proviso that if L is aminopiperazine, Ar^1 and Ar^2 are different or

if a is 1

L is a monovalent radical - NR^3R^4 , - SR^3 or - OR^3

where

R^3 and R^4 have one of the meanings of R^1 and R^2 or for -- NR^3R^4 , R^3 and R^4 can form a cyclic structure of the form (o)



where

U is an C_4 - C_6 alkyl residue optionally substituted by a substituent of formula Z and optionally interrupted by heteroatoms or heteroatom-containing groups such as -O-, - NR^1 ,

n is 1, 2 or 3 and

Z is hydrogen, optionally substituted C_1 - C_4 alkyl, - OR^5 , - CO_2R^5 , - COR^5

and

R^5 is hydrogen, optionally substituted C_1 - C_4 alkyl, optionally substituted vinyl, optionally substituted phenyl.

2. Dyestuff mixture according to claim 1

5 wherein


X^1 and X^2 is independently chlorine, fluorine or 3 or 4-carboxypyridinium;

Ar^1 and Ar^2 is independently a naphthyl residue substituted by at least one sulfo group (q-1)



where n is 1 to 3

or is a phenyl residue substituted by at least one sulfo group (q-2)



15 wherein

m is 1 or 2

p is 1 or 2 and

Y is independently hydrogen, halogen, R^5 , OR^5 , SR^5 , $NHCOR^5$,

where R^5 is as given in claim 1.

20


3. Dyestuff mixture according to claim 1

wherein

X^1 and X^2 is chlorine;


Ar^1 and Ar^2 are independently a naphthyl residue of the formula (q-11) or

25 (q-12)




(q-11)

or



(q-12)

or are a phenyl residue substituted by at least one sulfo group (q-2)



wherein

m is 1 or 2

p is 1 or 2 and


Y is methyl.

4. Dyestuff mixture according to claim 1 wherein a is 1 and L is morpholine.

5. A dye mixture according to at least one of the claims 1 to 4 wherein a dye of formula (I) is present in the mixture in an amount of from 1% by weight to 99% by weight and a dye of the formula (II) is present in the mixture in an amount of from 99% by weight to 1% by weight.

6. A dye mixture according to at least one of the claims 1 to 4 wherein a dye of formula (I) is present in the mixture in an amount of from 10% by weight to 90% by weight and a dye of the formula (II) is present in the mixture in an amount of from 90% by weight to 10% by weight.

7. A process for preparing a dye mixture as claimed in one or more of claims 1 to 5, which comprises reacting chromophores of formula (III)



(III)

wherein Ar¹, Ar², X¹, X² and M are as defined in claim 1 with an appropriate mixture of 2-aminoethylpiperazine and a diamine H-L-H, or amine H-L, wherein L is as defined above, followed by precipitation using methylated spirits and conventional filtration.

8. A process for dyeing hydroxy- and/or carboxamido – containing fiber material, in which dyestuffs or dyestuff mixtures are applied to the material and the dyes are fixed to the material by means of heat or with the aid of an alkali or by means of heat and with the aid of an alkali, which comprises dye mixtures or dyestuffs as claimed in one or more of the claims 1 to 5.

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C09B67/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C09B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 693 538 A (CIBA GEIGY AG) 24 January 1996 (1996-01-24) cited in the application claim 1	1-8
Y	US 5 892 006 A (TZIKAS ATHANASSIOS) 6 April 1999 (1999-04-06) claim 1	1-8
Y	DE 100 64 496 A (DYSTAR TEXTILFARBEN GMBH & CO) 4 July 2002 (2002-07-04) claim 1	1-8
A	WO 99 05244 A (CONNOR DANIEL STEDMAN ;SCHEIBEL JEFFREY JOHN (US); VINSON PHILLIP) 4 February 1999 (1999-02-04) cited in the application claim 1	1-8
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

25 February 2004

Date of mailing of the international search report

08/03/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bakboord, J

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 08104 A (BASF AG ;EBENEZER WARREN JAMES (GB); MYNETT DONNA MARIA (GB)) 17 February 2000 (2000-02-17) cited in the application claim 1	1-8
P,A	WO 02 092697 A (DYSTAR TEXTILFARBEN GMBH & CO ;EBENEZER WARREN JAMES (GB)) 21 November 2002 (2002-11-21) claim 1	1-8
P,A	EP 1 380 621 A (DYSTAR TEXTILFARBEN GMBH & CO) 14 January 2004 (2004-01-14) claim 1	1-8

Patent document cited in search report		Publication date		Patent family member(s)		Publication date	
EP 0693538		A		24-01-1996			
				BR		9502861 A	
				CN		1133323 A , B	
				DE		59509529 D1	
				EP		0693538 A2	
				ES		2161847 T3	
				GR		3036651 T3	
				JP		8003469 A	
				PT		693538 T	
				US		5631352 A	
US 5892006		A		06-04-1999			
				US		5735911 A	
				US		5612463 A	
				US		5451665 A	
				DE		59108779 D1	
				DE		59109196 D1	
				DE		59109201 D1	
				EP		0478503 A2	
				EP		0735107 A2	
				EP		0735113 A2	
				ES		2106070 T3	
				ES		2150620 T3	
				ES		2152455 T3	
				HK		1001093 A1	
				JP		3231364 B2	
				JP		4258674 A	
				JP		3389555 B2	
				JP		2001026726 A	
				JP		2002194241 A	
				US		5232462 A	
DE 10064496		A		04-07-2002			
				DE		10064496 A1	
				BR		0116419 A	
				CA		2428054 A1	
				WO		02051944 A1	
				EP		1345994 A1	
WO 9905244		A		04-02-1999			
				AU		738353 B2	
				AU		8124998 A	
				BR		9811524 A	
				CA		2297171 C	
				CN		1270622 T	
				EP		1002031 A1	
				HU		0002626 A2	
				WO		9905244 A1	
				ID		28301 A	
				JP		2001511474 T	
				TR		200000923 T2	
				US		2002103096 A1	
				US		6306817 B1	
				ZA		9806447 A	
WO 0008104		A		17-02-2000			
				AT		237661 T	
				BR		9912628 A	
				DE		69906983 D1	
				DE		69906983 T2	
				EP		1100847 A1	
				ES		2197658 T3	
				WO		0008104 A1	

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0008104	A		JP 2002522587 T	23-07-2002
			PT 1100847 T	31-07-2003
			TR 200100320 T2	21-06-2001
			US 6359121 B1	19-03-2002
WO 02092697	A	21-11-2002	CA 2446784 A1	21-11-2002
			WO 02092697 A1	21-11-2002
			EP 1387865 A1	11-02-2004
EP 1380621	A	14-01-2004	CA 2434825 A1	10-01-2004
			EP 1380621 A1	14-01-2004

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.